

Taking JIDEYTRO

JideytroTM
(zidesamtinib) | 25 mg
100 mg tablets

A guide to help you start and
stay on track with treatment

Patient portrayal.

What is JIDEYTROTM (zidesamtinib)?

JIDEYTRO is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that:

- has spread within the chest or other parts of the body and is caused by an abnormal ROS1 gene, and
- who have received a ROS1 kinase inhibitor.

It is not known if JIDEYTRO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

JIDEYTRO can cause serious side effects, including:

- **Central nervous system (CNS) effects.** Tell your healthcare provider right away if you or your family member taking JIDEYTRO develop any new or worsening CNS symptoms during treatment, including:
 - dizziness
 - fainting
 - walking, balance, or coordination problems
 - forgetfulness, confusion, attention or thinking problems
 - speaking problems
 - seizures
 - mood or personality changes, including anxiety, irritability
 - depression, including thoughts of suicide or dying
 - seeing or hearing things that are not real (hallucinations)

Please see full Important Safety Information on [pages 8–10](#)
and full [Prescribing Information](#) and [Patient Information](#) for JIDEYTRO.

Starting treatment with JIDEYTRO

Your doctor prescribed JIDEYTRO—now what? Here are a few things to keep in mind as you get started. Read on to learn more


JIDEYTRO (pronounced jih-DAY-troh) is a treatment designed to target your advanced *ROS1*+ non-small cell lung cancer (NSCLC)



Once-daily dosing



Can be taken with or without food



No sun exposure restrictions included in the full Prescribing Information

ROS1+ = ROS proto-oncogene 1-positive.

IMPORTANT SAFETY INFORMATION (cont.)

JIDEYTRO can cause serious side effects, including: (cont.)

- **Changes in the electrical activity of your heart called QT prolongation.** QT prolongation can cause irregular heartbeats that can be life-threatening or cause sudden death. Your healthcare provider will do tests before and during your treatment with JIDEYTRO to check the electrical activity of your heart and your body salts (electrolytes). Tell your healthcare provider right away if you feel faint, lightheaded, dizzy, or feel your heart beating irregularly or fast during your treatment with JIDEYTRO.

Please see full Important Safety Information on pages 8–10 and full Prescribing Information and Patient Information for JIDEYTRO.

What to know before taking JIDEYTRO

Before taking JIDEYTRO, tell your healthcare provider about all your medical conditions, including if you:

- Have nervous system (neurological) or psychiatric problems
- Have heart problems, including a condition called long QT syndrome
- Have lung or breathing problems other than lung cancer
- Are pregnant or plan to become pregnant. JIDEYTRO can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with JIDEYTRO

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with JIDEYTRO
- Use effective birth control (contraception) during treatment and for 6 months after the last dose of JIDEYTRO
- Talk to your healthcare provider about birth control methods that may be right for you

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment and for 3 months after the last dose of JIDEYTRO
- Are breastfeeding or plan to breastfeed. It is not known if JIDEYTRO passes into your breast milk. Do not breastfeed during treatment and for 1 week after your last dose of JIDEYTRO. Talk to your healthcare provider about the best way to feed your baby during treatment with JIDEYTRO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking JIDEYTRO with certain other medicines may affect the amount of JIDEYTRO or other medicines in your blood and may cause side effects or affect the way that JIDEYTRO or other medicines work.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

What should I avoid while taking JIDEYTRO?

Do not drive or operate heavy machinery during treatment if you develop side effects that can impact your mental alertness or coordination.

How should I take JIDEYTRO?



Take JIDEYTRO exactly as your healthcare provider tells you to take it. Do not change your dose or stop taking JIDEYTRO unless your healthcare provider tells you to



Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with JIDEYTRO if you develop side effects



Take JIDEYTRO at about the same time each day with or without food



Swallow JIDEYTRO tablets whole. Do not split, chew, crush, or dissolve the tablets. Do not take a tablet if it is broken, cracked, or damaged.



If you miss a dose of JIDEYTRO take it as soon as possible. If it has been 12 or more hours since you took your last dose, just skip the dose and take it at your next regularly scheduled time.



If you vomit at any time after taking JIDEYTRO, do not take another dose. Wait to take your next dose at your regularly scheduled time.

Make sure to store JIDEYTRO at **room temperature** between **68 °F to 77 °F** (20 °C to 25 °C).
Keep JIDEYTRO and all medicines out of the reach of children.

IMPORTANT SAFETY INFORMATION (cont.)

JIDEYTRO can cause serious side effects, including: (cont.)

- **Lung problems.** JIDEYTRO can cause lung problems that are severe or life-threatening. Tell your healthcare provider right away if you get any new or worsening symptoms of lung problems, including trouble breathing, shortness of breath, cough (with or without mucus), or fever.

Please see full Important Safety Information on [pages 8–10](#) and full [Prescribing Information](#) and [Patient Information](#) for JIDEYTRO.

Make JIDEYTRO a part of your day

Taking JIDEYTRO at the same time each day can help make treatment part of your routine

Creating this daily routine can help you:

- Remember to take your dose as prescribed
- Stay consistent with your treatment



Not actual size.



Helpful tips to consider:

- Set a daily alarm or reminder for yourself
- Keep JIDEYTRO in the same place
- Mark off your dose on a calendar or a phone app
- Ask a friend, family member, or care partner to help remind you

Your care team is here to help

Everyone's experience with treatment looks different. If you're having trouble remembering to take JIDEYTRO, talk with your healthcare provider. They can work with you to explore other strategies and tools that support your routine.

This information is not a substitute for medical advice. Always follow your doctor's instructions when taking JIDEYTRO.

JideytroTM
(zidesamtinib) | 25 mg
100 mg tablets

The safety and side effects of JIDEYTRO were studied in 446 people with ROS1+ cancer

Among the 446 were 432 people with ROS1+ NSCLC who received at least 1 dose of JIDEYTRO at 100 mg. They received JIDEYTRO once daily until their cancer progressed or they needed to stop treatment. Serious side effects occurred in 22% of these people. Serious side effects occurring in at least 2% of people were pneumonia and shortness of breath.

Most common side effects in 15% or more of people who received JIDEYTRO (out of 446):

Swelling (edema)	Numbness or tingling in arms or legs	Constipation	Tiredness	Shortness of breath
38%	25%	17%	16%	15%

The most common severe abnormal blood test results include increased creatine phosphokinase (CPK), increased triglycerides, decreased white blood cell counts and decreased red blood cell counts.

JIDEYTRO may cause fertility problems in males and females. Talk to your healthcare provider if fertility is a concern for you.

These are not all the possible side effects of JIDEYTRO. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088 (1-800-332-1088).

For more information, go to www.JIDEYTRO.com or call 1-844-NUVL-111 (1-844-688-5111).

Treatment experiences can vary—keep track of any side effects and discuss them with your care team.

IMPORTANT SAFETY INFORMATION (cont.)

JIDEYTRO can cause serious side effects, including: (cont.)

- **Bone fractures.** JIDEYTRO can increase your risk of bone fractures. Bone fractures may happen with or without a fall or other injury. Tell your healthcare provider if you develop pain, changes in movement, or bone abnormalities.

Please see full Important Safety Information on pages 8–10 and full Prescribing Information and Patient Information for JIDEYTRO.

Tolerability that may help you stay on therapy

Most side effects did not require people to permanently stop treatment or lower their dose in the JIDEYTRO clinical study

2.3%

(10 out of 432 people) permanently stopped treatment because of side effects.

10%

(43 out of 432 people) needed their dose reduced because of side effects.

The most common reasons for stopping treatment included pneumonia and pneumonitis.

Treatment experiences can vary—keep track of any side effects and discuss them with your care team.

IMPORTANT SAFETY INFORMATION

What is JIDEYTRO™ (zidesamtinib)?


JIDEYTRO is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that:

- has spread within the chest or other parts of the body and is caused by an abnormal ROS1 gene, and
- who have received a ROS1 kinase inhibitor.

It is not known if JIDEYTRO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

JIDEYTRO can cause serious side effects, including:




Central nervous system (CNS) effects. Tell your healthcare provider right away if you or your family member taking JIDEYTRO develop any new or worsening CNS symptoms during treatment, including:

- dizziness
- fainting
- walking, balance, or coordination problems
- forgetfulness, confusion, attention or thinking problems
- speaking problems
- seizures
- mood or personality changes, including anxiety, irritability
- depression, including thoughts of suicide or dying
- seeing or hearing things that are not real (hallucinations)




Changes in the electrical activity of your heart called QT prolongation. QT prolongation can cause irregular heartbeats that can be life-threatening or cause sudden death. Your healthcare provider will do tests before and during your treatment with JIDEYTRO to check the electrical activity of your heart and your body salts (electrolytes). Tell your healthcare provider right away if you feel faint, lightheaded, dizzy, or feel your heart beating irregularly or fast during your treatment with JIDEYTRO.



Lung problems. JIDEYTRO can cause lung problems that are severe or life-threatening. Tell your healthcare provider right away if you get any new or worsening symptoms of lung problems, including trouble breathing, shortness of breath, cough (with or without mucus), or fever.



Bone fractures. JIDEYTRO can increase your risk of bone fractures. Bone fractures may happen with or without a fall or other injury. Tell your healthcare provider if you develop pain, changes in movement, or bone abnormalities.



Muscle pain, tenderness, and weakness (myalgia). JIDEYTRO can cause myalgia with an increase in the level of an enzyme in your blood called creatine phosphokinase (CPK), which may be a sign of muscle damage. Your healthcare provider will do blood tests to check your CPK blood levels every 2 weeks during the first month, then every 1 to 2 months and as needed if you get unexplained muscle pain, tenderness, or weakness during your treatment with JIDEYTRO. Tell your healthcare provider if you develop any of these symptoms.



Inflammation of the pancreas (pancreatitis). JIDEYTRO may increase enzymes in your blood called amylase and lipase, which may be a sign of pancreatitis. Your healthcare provider will do blood tests to check your pancreatic enzyme blood levels during treatment with JIDEYTRO. Tell your healthcare provider right away if you get any signs and symptoms of pancreatitis, including upper stomach (abdominal) pain that may spread to the back and get worse with eating, weight loss, or nausea.

Before taking JIDEYTRO, tell your healthcare provider about all your medical conditions, including if you:

- have nervous system (neurological) or psychiatric problems.
- have heart problems, including a condition called long QT syndrome.
- have lung or breathing problems other than lung cancer.
- are pregnant or plan to become pregnant. JIDEYTRO can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with JIDEYTRO.

Females who are able to become pregnant: Your healthcare provider will do a pregnancy test before you start treatment with JIDEYTRO. Use effective birth control (contraception) during treatment and for 6 months after the last dose of JIDEYTRO. Talk to your healthcare provider about birth control methods that may be right for you.

Males with a female partner who are able to become pregnant: Use effective birth control during treatment and for 3 months after the last dose of JIDEYTRO.

IMPORTANT SAFETY INFORMATION (cont.)

Before taking JIDEYTRO, tell your healthcare provider about all your medical conditions, including if you: (cont.)

- are breastfeeding or plan to breastfeed. It is not known if JIDEYTRO passes into your breast milk. **Do not** breastfeed during treatment and for 1 week after your last dose of JIDEYTRO. Talk to your healthcare provider about the best way to feed your baby during treatment with JIDEYTRO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking JIDEYTRO with certain other medicines may affect the amount of JIDEYTRO or other medicines in your blood and may cause side effects or affect the way that JIDEYTRO or other medicines work.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

What should I avoid while taking JIDEYTRO?

- Do not drive or operate heavy machinery during treatment if you develop side effects that can impact your mental alertness or coordination.

What are the most common side effects of JIDEYTRO?

The most common side effects of JIDEYTRO include:

- swelling (edema)
- tiredness
- numbness or tingling in your arms or legs
- shortness of breath
- constipation

The most common severe abnormal blood test results include increased creatine phosphokinase (CPK), increased triglycerides, decreased white blood cell counts and decreased red blood cell counts.

JIDEYTRO may cause fertility problems in males and females. Talk to your healthcare provider if fertility is a concern for you.

These are not all the possible side effects of JIDEYTRO. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see full [Prescribing information](#) and [Patient Information](#) for JIDEYTRO.



JideytroTM
(zidesamtinib) | 25 mg
100 mg tablets

Guiding a path to treatment: Nuvalent Patient Support

We're here to help you understand what to expect with your medication, navigate insurance coverage and costs, and start and stay on JIDEYTRO (zidesamtinib) with the following programs:

Patient Assistance Program*

If you do not have insurance, your insurance does not cover JIDEYTRO, or you are concerned about the cost of your medication even with insurance, the Patient Assistance Program may be able to provide your medication at no cost if you meet eligibility requirements.

Quick Start Program*

If you are new to JIDEYTRO and there is a delay in insurance review or coverage, you may be eligible to receive a free, temporary supply of medication while you wait for a coverage decision.

Bridge Program*

If you are already taking JIDEYTRO and your insurance changes, you may be eligible for a free, temporary supply of medication while you wait for new insurance or coverage.

Benefits Verification† and Coverage Information

Our team can help you and your healthcare provider understand insurance coverage for your medication, the steps needed to receive approval from your plan, and what costs you may be responsible for.

Copay Assistance Program

You could pay as little as \$0 for JIDEYTRO if you have commercial insurance. Patients covered by Medicare, Medicaid, or other state/federal plans are not eligible. Maximum benefit per calendar year applies.

Advocacy and Community Groups

Our team can connect you to patient advocacy organizations that offer support and resources for patients and their care partners that may be tailored to your specific needs.

Start exploring advocacy organizations on the back page.

Talk to your healthcare provider or visit our website to enroll in the above programs.‡

Enroll in Nuvalent Patient Support
NuvalentPatientSupport.com

*Please see the Patient Assistance Program, Quick Start Program, and Bridge Program terms and conditions.

†Verification of benefits is not a guarantee of payment and does not take the place of written policy information. Healthcare providers should carry out their own benefits investigation as necessary.

‡Please see the Copay Assistance Program terms and conditions.



Patient portrayal.

Jideytro™
(zidesamtinib) | 25 mg
100 mg tablets

Connect to others living with ROS1+ NSCLC

JideytroTM
(zidesamtinib) | 25 mg
100 mg tablets

Discover support for people with ROS1+ NSCLC
through these patient advocacy organizations:

The **ROS1**ders

GO2
FOR LUNG
CANCER

**LUNG CANCER
RESEARCH
FOUNDATION**

LCFA
Lung Cancer Foundation of America

LUNGEVITY
Transforming Lung Cancer

Patient portrayal.

These organizations are separate from Nuvalent and are not connected to JIDEYTRO. They are listed to provide extra information and support, and they should not replace medical advice from your care team.

IMPORTANT SAFETY INFORMATION (cont.)

JIDEYTRO can cause serious side effects, including: (cont.)

- **Muscle pain, tenderness, and weakness (myalgia).** JIDEYTRO can cause myalgia with an increase in the level of an enzyme in your blood called creatine phosphokinase (CPK), which may be a sign of muscle damage. Your healthcare provider will do blood tests to check your CPK blood levels every 2 weeks during the first month, then every 1 to 2 months and as needed if you get unexplained muscle pain, tenderness, or weakness during your treatment with JIDEYTRO. Tell your healthcare provider if you develop any of these symptoms.

**Please see full Important Safety Information on pages 8–10
and full Prescribing Information and Patient Information for JIDEYTRO.**



Nuvalent and the associated logo are registered trademarks of Nuvalent, Inc., and JIDEYTRO, Nuvalent Patient Support, and the associated logos are trademarks of Nuvalent, Inc.

© 2026 Nuvalent. All rights reserved. ZID-US-0044v1.0 07/2026